

Change of phytoalexins -stilbenoids of vine leave Tsitska variety (*Vitis vinifera* L.) in condition Downy mildew

M. Bezhuashvili ^{1*}, L. Tskhvedadze ¹, M. Surguladze ¹, G. Shoshiashvili ¹, L. Elanidze ¹, P. Vashakidze ¹

¹ Institute of Viticulture and Oenology of Agricultural University of Georgia, Tbilisi, GEORGIA

*Corresponding author: m.bezhuashvili@agruni.edu.ge

Abstract

It is investigated change of phytoalexin-stilbenoids in the leaves of white grapevine Tsitska infected by Downy mildew. The Tsitska is cultivated in the west part of Georgia. Comparing the healthy and diseased grapevine leaves physiological concentrations of stilbenoids, in the infected grapevine leaves it is established following stress-metabolite stilbenoids: trans-resveratrol and ϵ -viniferin. The stress-metabolite stilbenoids in the conditions of the disease with Downy mildew is a scientific novelty for the vine leave Tsitska variety. The results of the research are important for determination the correlation of the vine immunity with the phytoalexins-stilbenoids.

Keywords: phytoalexin-stilbenoids, Tsitska, Downy mildew

Bezhuashvili M, Tskhvedadze L, Surguladze M, Shoshiashvili G, Elanidze L, Vashakidze P (2020) Change of phytoalexins -stilbenoids of vine leave Tsitska variety (*Vitis vinifera* L.) in condition Downy mildew. Eurasia J Biosci 14: 167-171.

© 2020 Bezhuashvili et al.

This is an open-access article distributed under the terms of the Creative Commons Attribution License.

INTRODUCTION

Vine and grape stilbenoids are one of the groups of a wide class of phenol compounds, which incorporates cis- and trans-isomers of monomer resveratrol and their derivatives, as dimers, trimers, tetramers and glycosides (Aaviksaar et al. 2003, Bavaresco et al. 2002, Guebailia et al. 2006, Larronde et al. 2005, Niesen Daniel 2013, Pawlus et al. 2012, Reniero et al. 1996, Rivière et al. 2012, Waffo Teguo et al. 1998) Stilbenoids have diversified high biological activity and these compounds are very important for plants, as phytoalexins. Phytoalexin characteristics of grape vine is established by us in "in vitro" experiments by their inhibitory impact on the microorganisms, which are due to grape vine bacterial and fungus diseases. These diseases are: Grey mildew (*Botrytis cinerea*), Downy mildew (*Plasmopara viticola*), Powdery mildew (*Uncinula necator*), crown gall (*Agrobacterium tumefaciens*) (Bezhuashvili et al. 2019a, 2019b). Stilbenoids act against different vine diseases caused by biotic factors. The following stilbenoids were identified in the extract of vine (*Vitis vinifera*) trunk, roots and annual shoots: Ampelopsin A, (E)-piceatannol, Pallidol, E-resveratrol, hopeaphenol, isohopeaphenol, (E)- ϵ -viniferin, (E)-miyabenol C, (E)-w-viniferin, r- and r2-viniferin. It was established that the extract inhibits the growth of sporulation of fungus *Plasmopara viticola* by 50%, while the most active inhibitor of it turned out to be r2-viniferin (Gabaston et al. 2017). Under the influence of *Botrytis cinerea* on the mixture of Pterostilben and

Resveratrol 7 new stilbens were formed, while 5 new stilbens were formed from Pterostilben under the same terms. The anti-fungus effect of these stilbenoids was fixed against *Plasmopara viticola* (Gindro et al. 2017). At three stages of the grape (*Vitis vinifera*) grain development, the grains were infected on purpose with *Botrytis cinerea* spores "in vitro." In the infected grain, stilbenoids: Pterostilben, (E)- ϵ -viniferin and trans-resveratrol were fixed. Dominating among them was (E)- ϵ -viniferin (Bavaresco et al. 1997). The grains of *Vitis Vinifera* L. cv. Barbera in the ripening period were infected with conidial suspension of *Aspergillus jannicus*, *A.ochraceus*, *A. fumigatus* and *A.carbonariuces*. The process of formation of ochratoxin A and stilbenoids was supervised. It was found out that all experimental fungi except *A. Fumigatus* significantly increase the concentration of trans-resveratrol and at the same time, trans-Piceid stays unchanged. In the grape grain damaged by *A.ochraceus*, the concentration of piceatannol increased significantly. A large amount of *A.carbonariuce* was synthesized in the grain infected with *A.carbonariuces* isolate and the anti-fungicidal activity occurred with the following concentrations: 300 mkg gr⁻¹ and 20 mkg gr⁻¹, what was sufficient for the total inhibition of fungus *A.carbonariuces* (Bavaresco et al. 2003). Besides above mentioned biological activity

Received: April 2019

Accepted: November 2019

Printed: February 2020




Fig. 1. Chart of isolating a stilbenoid-containing fraction from vine leaf

stilbenoids have many other functional purposes (Bavaresco and Fregoni 2001, Bavaresco et al. 2003, Gabastoni et al. 2018, Jeandet et al. 2002, Langcake and Pryce 1976, Langcake et al. 1979, Langcake 1981, Mattivi et al. 2011, Sáeza et al. 2018). The vine and grape impacts some factor (Adrian et al. 2000, Bavaresco 2003, Bavaresco et al. 2007, Vergara et al. 2012). The vine varieties of Georgia are rich in biologically active stilbenoids. *trans-resveratrol*, *trans-ε-viniferin*, 2 tetrameric stilbens, including hopeaphenol as one of them, were isolated and identified from an annual shoot of *Rkatsiteli* variety. These stilbenoids and new stilbenoids identified by us were identified in the Georgian red-grape vintage varieties and their wines (Bezhuashvili 1994, Bezhuashvili and Surguladze 2016, Bezhuashvili et al. 2013, Surguladze and Bezhuashvili 2017, 2018). The study of stilbenoids in Georgian vintage varieties as that of phytoalexins, qualitative and quantitative analyses of their physiological concentrations and stress-metabolites and their impact on the microorganisms causing bacterial and fungus diseases is an urgent issue of the research. Consequently, our goal was to identify the vine varieties

infected with crown gall disease, identify and determine their stress-metabolite stilbenoids and compare them with healthy vine stilbenoid profile. It is established change of stilbenoids healthy and infected (95% and 50%) vine leaves and canes from 35 years old vineyard of cabernet franc (france). It is identified *E-piceatannol*, *E-resveratrol*, *E-ε-viniferin*, *ampelopsin A*, *E-miyabenol C*, *E-vitisin B*, *hopeaphenol*, *isohopeaphenol*. In infected grape skin was identified high quotation of *E-ε-viniferin* to compare with *E-resveratrol*. To point of view of authors: "These findings suppose that the health status in vineyards could modify the composition of stilbenoids in winter-harvested grape canes and subsequently the potential biological properties of the valuable extract (Houlline et al. 2015).

Research Aim

Research aim was to establish vine sort of *tsitska* leaves correlation with phytoalexin stilbenoid change infected by downy mildew.




Fig. 2. Change of concentration (mg/kg) of trans-resveratrol and trans-epsilon-viniferin in condition Downy mildew in the leaves of Tsitska variety

MATERIALS AND METHODS

Research objects were healthy and infected by downy mildew leaves of Georgian white grape tsitska in west part of Georgia (Zestafoni region). Vineyards are located raw humus calcareous-rendzic-leptosols type of soil 30 years of vine. We isolated stilbenoid-containing fractions from the healthy and infected vine leave as a result of treatment according to the chart (**Fig. 1**).

Stilbenoids were determined by the method of high-performance liquid chromatography (HPLC) (Guebailia et al. 2006). For this purpose, we used the Varion chromatograph SupelcosilPM LC18 Column, 250x4,6mm, eluents: A. 0.025% trifluoroacetic acid, B.Acetonitrile: A80/20. Gradient mode: 0-35 min, 20-

50% B, 48-53min, 200% B. Flow rate of the eluent- 1 ml/min; wavelength-306 and 285nm. The samples were analyzed three times and it is presented average results. Analyzed samples: isolated stilbenoid-containing fractions were filtered using a membrane filter (0,45 μ) before the chromatographic procedure. The chromat-mass-spectral investigations were carried out under the above-mentioned conditions; mass-spectra were detected by obtaining of positive ions.

RESULTS AND DISCUSSION

During the experiment it was determined quantitative change of phytoalexin-stilbenoids in condition of infected by Downy mildew vine leaves. Depending on the received results established main stress-metabolite stilbenoids: trans-resveratrol and trans- ϵ -viniferin. As diagram presents the physiological concentration of both compounds are increasing in condition downy mildew infect,in concret: trans-resveratrol 0.82 mg/kg-19.32 mg/kg and trans- ϵ -viniferin 8.56mg/kg-10.25mg/kg.

CONCLUSION

Established first researched results is scientific novelty for vine leave Tsitska variety. Belong on the basic of this and future research results will be found out the correlation of vine Tsitka variety with phytoalexins - stilbenoids.

This work was supported by Shota Rustaveli National Science Foundation of Georgia (SRNSFG) [grant number FR 17_486].

REFERENCES

- Aaviksaar A, Haga M, Püssa T, Roasto M, Tsoupras G (2003) Purification of resveratrol from vine stems. Proc. Estonian Acad. Sci. Chem 54: 155-164.
- Adrian M, Jeandet P, Douillet-Breuil A, Tesson L, Bessis R (2000) Stilbene content of mature *Vitis vinifera* berries in response to UV-C elicitation. J. Agric. Food Chem 48: 6103-6105.
- Bavaresco L (2003) Role of viticultural factors on stilbene concentrations of grapes and wine. Drugs Exp. Clin. Res 29: 181-187.
- Bavaresco L, Fregoni C (2001) Physiological role and molecular aspects of grapevine stilbenic compounds, Molecular biology & biotechnology of the grapevine. Kluwer Academic Publishers, The Netherlands 500: 153-182.
- Bavaresco L, Fregoni M, Trevisan M, Mattivi F, Vrhovsek U, Falchetti R (2002) The occurrence of the stilbene piceatannol in grapes. Vitis 41: 133-136.
- Bavaresco L, Petegolli D, Cantù E, Fregoni M, Chiusa G, Trevisan M (1997) Elicitation and accumulation of stilbene phytoalexins in grapevine berries infected by *Botrytis cinerea*. Vitis: 3677-3683.
- Bavaresco L, Pezzutto S, Fornaroli A, Ferrari F (2003) Grapevine iron-chlorosis occurrence and stilbene root concentration as affected by the rootstock and arbuscular mycorrhizal infection L. Acta Hort 603: 401-410.
- Bavaresco L, Pezzutto S, Gatti M, Mattivi F (2007) Role of the variety and some environmental factors on grape stilbenes. Vitis 46: 57-61.
- Bavaresco L, Vezzulli S, Battilani P, Giorni P, Pietri A, Bertuzzi T (2003) Effect of Ochratoxin A-Producing *Aspergilli* on Stilbenic Phytoalexin Synthesis in Grapes. J.Agric.Food.Chem 51: 6151-6157. <https://doi.org/10.1021/jf0301908>

- Bezhuashvili M (1994) Development of theoretical basics for vine and oak timber lignin and identification of the ways to use the gained products. Doctoral thesis, Tbilisi.
- Bezhuashvili M, Surguladze M (2016) The Derivatives of Resveratrol-Viniferins from Grape Saperavi (*Vitis vinifera* L.). 3rd Scientific Conference on Natural and Synthetic Biological Activity Substances, Georgian National Academy of Sciences, TSMU I. Kutateladze Institute of Pharmacochimistry. 24-25 October, Tbilisi, Georgia. Abstract Book: 112-113.
- Bezhuashvili M, Tskhvedadze L, Surguladze M, Shoshiashvili G, Gagunashvili L, Elanidze L, Vashakidze P (2019) Change of Phytoalexin Stilbenoids of Grape Skin Rkatsiteli (*Vitis vinifera* L.) in condition of Powdery Mildew, Bioscience Biotechnology Research communications 12(5): 146-151.
- Bezhuashvili M, Tskhvedadze L, Surguladze M, Shoshiashvili G, Kharadze SH, Gagunashvili L, Elanidze L, Vashakidze P (2019) Stress-metabolites phytoalexins -stilbenoids of grape skin Rkatsiteli variety (*Vitis vinifera* L.) in condition gray mildew. Euroasian Journal of Biosciences 13: 549-554.
- Bezhuashvili, M., Shubladze, L., Okruashvili, D. (2013). Trans-Piceid Stilbenoid in the Juice and Skin of the red Grape (*Vitis vinifera* L.) Varieties growing in Georgia. Bull.Georg.Nati.Acad.Sci 7(3): 774-779.
- Gabaston J, Cantos-Villar E, Biais B, Waffo-Teguo P, Renouf E, Corio-Costet MF, Richard T, Mérillon JM (2017) Stilbenes from *Vitis vinifera* L. Waste: A Sustainable Tool for Controlling Plasmopara Viticola. J.Agric. Food Chem 65: 2711-2718.
- Gabastoni J, Khawand E, Waffo-Teguo P, Decendit A, Richard T, Mérillon JM, Pavela R (2018) Stilbenes from grapevine root: a promising natural insecticide against *Leptinotarsa decemlineata*. Journal of Pest Science 91(2): 897-906. <https://doi.org/10.1021/jf303290g>
- Gindro K, Schnee S, Righi D, Marcourt L, Ebrahimi SN, Codina M, Voinesco F, Michellod E, Wolfender J-L, Queiroz EF (2017) Generation of Antifungal Stilbenes Using the Enzymatic Secretome of *Botrytis cinerea*. Journal of Natural products 80: 887-898.
- Guebailia HA, Chira K, Richard T, Mabrouk T, Furiga A (2006) Hopeaphenol: the first resveratrol tetramer in wines from North Africa. J.Agric. Food Chem 54: 9559-9564. <https://doi.org/10.1021/jf062024g>
- Houlline B, Besseau S, Delanoue G, Oudin A, Papon N, Giglioli-Guivarch M, Zanoué A (2015) Composition and Tissue-Specific Distribution of Stilbenoids in Grape Canes are affected by Downy Mildew pressure in the vineyard. J.Agricultural and Food chemistry 63: 8472-8477. <https://doi.org/10.1021/acs.jafc.sbo2997>
- Jeandet, P., Bessis, R., Douillet-Breuil, A., Adrian, M. (2002). Phytoalexins from the Vitaceae: Biosynthesis, Phytoalexin Gene Expression in Transgenic Plants, Antifungal Activity, and Metabolism. Article · Literature Review in Journal of Agricultural and Food Chemistry 50(10): 2731-2741. <https://doi.org/10.1021/jf950807o>
- Langcake P (1981). Disease resistance of *Vitis* spp. and the production of the stress metabolites resveratrol, epsilon-viniferin, alpha-viniferin and pterostilbene. Physiol. Plant Pathol 18: 213-226.
- Langcake P, Cornford CA, Pryce RJ (1979) Identification of pterostilbene as a phytoalexin from *Vitis vinifera* leaves. Phytochemistry 18: 1025-1027. [https://doi.org/10.1016/S0031-9422\(00\)91470-5](https://doi.org/10.1016/S0031-9422(00)91470-5)
- Langcake P, Pryce RJ (1976) The production of resveratrol by *Vitis vinifera* and other members of the vitaceae as a response to infection or injury. Physiol. Plant Pathol: 977-86
- Larronde F, Richard T, Delaunay JC, Decendit A, Monti JP, Krisa S, Mérillon JM (2005) New stilbenoid glucosides isolated from *Vitis vinifera* cell suspension cultures (cv. Cabernet sauvignon). Planta Medica 71: 888-890.
- Mattivi F, Vrhovsek U, Malacarne G, Masuero D, Luca Zulini Stefanini M, Moser C, Velasco G, Guella R (2011) Profiling of Resveratrol Oligomers, Important Stress Metabolites, Accumulating in the Leaves of Hybrid *Vitis vinifera* (Merzling × Teroldego) Genotypes Infected with *Plasmopara viticola*. J.Agric.FoodChem.59,5364-5375.
- Niesen Daniel B, Hessler C, Navindra SP (2013) Beyond resveratrol: A review of natural stilbenoids identified from 2009-2013. Journal of Berry Research 3: 181-196.
- Pawlus D, Waffo-Teguo P, Shaver J, Merillon JM (2012) Stilbenoid Chemistry from Wine and the Genus *Vitis*. Int. Sci. Vigne Vin 46(2): 57-111.
- Reniero F, Angioni A, Rudolph M, Mattivi F (1996) Identification of two stilbenoids from *Vitis* roots. Vitis 35: 125-127.
- Rivière C, Pawlus A, Merillon J-M (2012) Natural stilbenoids: distribution in the plant kingdom and chemotaxonomic interest in Vitaceae. Natural Product Reports 29(11): 1317-1333. <https://doi.org/10.1039/c2np20049>
- Sáeza V, Pasteneb E, Vergaraa C, Mardonesa C, Hermosín-Gutiérrezc I, Gómez-Alonsoc S, Gómezc V, Theodulozd C, Riquel Mee S, Baer D (2018) Oligostilbenoids in *Vitis vinifera* L. Pinot Noir grape cane extract: Isolation, characterization, in vitro antioxidant capacity and anti-proliferative effect on cancer cells panel. Food Chemistry 265: 101-110. <https://doi.org/10.1016/j.foodchem.2018.05.050>

- Surguladze M, Bezhuashvi M (2017) New stilbenoids - Miyabenols from Saperavi grapes (*Vitis vinifera* L.) to wine. 8 th international conference on polyphenols and health. Octobre 3-6, Quebec, Canada, Abstract Book: p. 52.
- Surguladze M, Bezhuashvili M (2018) Biological activity of some stilbenoids from grape saperavi (*Vitis vinifera* L.) ICP+TC 2018 XXIX International Conference on Polyphenols 9th Tannin Conference: 16-20. Madison, USA. [Conferences.union.wisc.edu/icp/abstracts/poster-presentations](https://conferences.union.wisc.edu/icp/abstracts/poster-presentations).
- Vergara C, von Baer D, Mardones C, Wilkens A, Wernekinck K, Damm A, Macke S, Gorena T, Winterhalter P (2012) Stilbene Levels in Grape Cane of Different Cultivars in Southern Chile: Determination by HPLC-DAD-MS/MS Method, *Agric. Food Chem* 60: 929-933.
- Waffo Teguo P, Fauconneau B, Deffieux G, Huguet F, Vercauteren J, Merillon JM (1998) Isolation, identification, and antioxidant activity of three stilbene glucosides newly extracted from *vitis vinifera* cell cultures. *J. Nat. Prod* 61: 655-657.

www.ejobios.org